

بِهِ نَاهِ خَدَا

دکتر اشرف آل یاسین

استاد گروه ذنان و ذایمان
دانشگاه علوم پزشکی تهران



Ovarian Rejuvenation

In poor responder & POI

1. POR 8-24%
2. POI (Occult, Biochemical, Overt POI)

Techniques rejuvenation

1. Activity Hippo signaling pathway IOA
2. Mitochondrial transfer pro cell (exchange)



Growth factors can improve tissue perfusion via intra stromal angiogenesis
oocyte quality improve intra follicula oxygen, recovery of mitochondrial
function

- 1) Only PRP
- 2) PRR + stem cell
- 3) PRP + gonadotropin

Success rate → dose dependent

Long lasting effect 14 month

Autologous stem cell therapy

1. Bone marrow
2. Ovary
3. Adipose tissue

Exact mechanism: unknown

- 1) Homing phenomenon
- 2) Para crin action
- 3) Differentiate into variety of cells
- 4) Promote angiogenesis
- 5) Reducing apoptosis
- 6) Primordial follicles ↑



Complication

Ovarian granulosa cell
tumor

A photograph of a field of red poppies growing among green grass and weeds. The poppies are in various stages of bloom, with some fully open and others still buds. The background is a soft-focus view of more flowers and plants. A thin blue circular border surrounds the image.

Thank you